


1. In order to balance the following chemical reaction



set up a system of linear equations for the variables x_i .

/10

- (a) Write the augmented matrix representing this linear system.

✓ solution

$$\left(\begin{array}{cccc|c} 10 & 0 & -1 & 0 & 0 \\ 16 & 0 & 0 & -1 & 0 \\ 0 & 2 & 0 & -1 & 0 \end{array} \right)$$

- (b) Find the solution set of the system.

✓ solution

$$\mathbf{x} = \alpha \begin{pmatrix} 1/16 \\ 1/2 \\ 10/16 \\ 1 \end{pmatrix}$$

- (c) Specialize the solution so that the values of the variables x_i are positive integers.

✓ solution

$$\text{For } \alpha = 16 \text{ the solution is } \mathbf{x} = \begin{pmatrix} 1 \\ 8 \\ 10 \\ 16 \end{pmatrix}.$$

2. Given a system of linear equations

$$\begin{aligned} x + y &= c_1 \\ 2x + 3y &= c_2 \\ x - y &= c_3 \end{aligned}$$

/15

- (a) Reduce the system to row-echelon form.

✓ solution

$$\left(\begin{array}{cc|c} 1 & 1 & c_1 \\ 0 & 1 & c_2 - 2c_1 \\ 0 & 0 & -5c_1 + 2c_2 + c_3 \end{array} \right)$$

- (b) For what values does the system have a solution? Explain.

✓ solution

Only if $-5c_1 + 2c_2 + c_3 = 0$ the system is consistent.

- (c) Find the best approximate solution for $c_1 = 1$, $c_2 = 2$, $c_3 = 1$.

✓ solution

$$\mathbf{x} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

(Note: since the given values satisfy the equation in (b), the best approximate solution is the exact solution)

3. Find a basis of the subspace of \mathbb{R}^4 of vectors orthogonal to $(1, 3, -1, 0)^T$ and $(1, 3, 0, 1)^T$.

/10

$$\left[\begin{pmatrix} -3 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \\ -1 \\ 1 \end{pmatrix} \right]$$

✓ solution

4. Given a matrix

$$A = \begin{pmatrix} 1 & 2 & 2 \\ 0 & 1 & 1 \\ 0 & -2 & 4 \end{pmatrix},$$

/10

Find the eigenvalues of A .

$$\lambda_1 = 1, \lambda_2 = 2 \text{ and } \lambda_3 = 3$$

✓ solution

5. Consider $\mathbf{p} = (3, 2, 1)$ and the plane $x + 2y + 2z = 0$.

/10

- (a) Find a unit normal vector to the plane.

$$\mathbf{n} = \frac{1}{3} \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$$

✓ solution

- (b) Find the distance from the point \mathbf{p} to the plane.

$$d = 3$$

✓ solution

- (c) Find the projection of \mathbf{p} onto the plane.

$$\mathbf{p} - d\mathbf{n} = \begin{pmatrix} 2 \\ 0 \\ -1 \end{pmatrix}$$

✓ solution

6. Find the symmetric matrix representing the quadratic form

$$q(x, y) = x^2 - 3y^2 + 2z^2 + 2xy - 2xz + 4yz.$$

/10

$$\begin{pmatrix} 1 & 1 & -1 \\ 1 & -3 & 2 \\ -1 & 2 & 2 \end{pmatrix}$$

✓ solution

7. Let $L_1 : P_3 \rightarrow P_3$ and $L_2 : P_3 \rightarrow P_3$ be the linear operators defined by

$$L_1(p) = p'' + 2p', \quad L_2(p) = p''(0) + 2p'(0),$$

where $p(x) \in P_3$.

/10

(a) Find the matrix representing L_1 in the standard basis $[1, x, x^2]$.

✓ solution

$$\begin{pmatrix} 0 & 2 & 2 \\ 0 & 0 & 4 \\ 0 & 0 & 0 \end{pmatrix}$$

(b) Find the matrix representing L_2 in the standard basis $[1, x, x^2]$.

✓ solution

$$\begin{pmatrix} 0 & 2 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

8. Use Gram-Schmidt orthogonalization to find an orthonormal basis for the space spanned by the vectors

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \\ 1 \end{pmatrix}, \quad \mathbf{v}_2 = \begin{pmatrix} 2 \\ 1 \\ 0 \\ 1 \end{pmatrix}, \quad \text{and} \quad \mathbf{v}_3 = \begin{pmatrix} 4 \\ 3 \\ -2 \\ 1 \end{pmatrix}.$$

/10

✓ solution

$$\left[\frac{1}{\sqrt{3}} \begin{pmatrix} 1 \\ 0 \\ 1 \\ 1 \end{pmatrix}, \frac{1}{\sqrt{3}} \begin{pmatrix} 1 \\ 1 \\ -1 \\ 0 \end{pmatrix} \right]$$

9. Let $A = \begin{pmatrix} 1 & 1 & -1 \\ 0 & 2 & 0 \\ 0 & 0 & -1 \end{pmatrix}$

/15

(a) Give all eigenvalues and corresponding eigenvectors of A .

✓ solution

$$\lambda_1 = 1, \quad \mathbf{v}_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

$$\lambda_1 = 2, \mathbf{v}_1 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$$
$$\lambda_1 = -1, \mathbf{v}_1 = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}$$

(b) Give the general form of the solution to $\mathbf{x}'(t) = A\mathbf{x}(t)$.

$$\mathbf{x}(t) = c_1 e^t \mathbf{v}_1 + c_2 e^{2t} \mathbf{v}_2 + c_3 e^{-t} \mathbf{v}_3$$

✓ solution

(c) If $\mathbf{x}(0) = \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix}$, what is then the solution to $\mathbf{x}'(t) = A\mathbf{x}(t)$?

$$c_1 = c_2 = 1, c_3 = -1$$

✓ solution

$$\mathbf{x} = \begin{pmatrix} e^t + e^{2t} - e^{-t} \\ e^{2t} \\ -2e^{-t} \end{pmatrix}$$